Example Report

ARUP Laboratories 500 Chipeta Way – Salt Lake City, UT 84108 (800)522-2787 - www.aruplab.com Julio C. Delgado, M.D. M.S., Director of Laboratories

Patient Age/Gender: Unknown Unknown Printed: 05-Dec-18 11:30:06

 $\frac{\text{Procedure}}{\text{Sulfonamides (Sulfas)}} = \frac{\text{Result}}{11.3} = \frac{\text{Units}}{\text{mg/dL}} = \frac{\text{Ref Interval}}{[5.0-15.0]} = \frac{\text{Accession}}{18-339-900050} = \frac{\text{Collected Received}}{05-\text{Dec}-18} = \frac{\text{Verified Note of the Note of the$

05-Dec-18 11:02:00 Sulfonamides (Sulfas): INTERPRETIVE INFORMATION: Sulfonamides (Sulfas)

Therapeutic Range: 5.0-15.0~mg/dL Toxic: Greater than 20.0~mg/dL

This test is designed to measure sulfamethoxazole. Peak sulfonamide (total) blood levels of 5.0--15.0 mg/dL may be effective for most infections, with concentrations of 12.0--15.0 mg/dL being optimal for serious infections. Sulfonamide levels should not exceed 20.0 mg/dL. Adverse effects may include blood dyscrasias, skin rash, nausea, vomiting and fever.

Chart ID: 13157356 Page 1 of 1