



<u>Procedure</u>	<u>Result</u>	<u>Units</u>	<u>Ref Interval</u>	<u>Accession</u>	<u>Collected</u>	<u>Received</u>	<u>Reported/ Verified</u>
HSV Type 1/2 Combined Ab, IgG	2.52 f	IV		19-344-900041	10-Dec-19 09:57:00	10-Dec-19 09:57:00	10-Dec-19 09:59:06
HSV 1 and/or 2 Abs, IgM by ELISA	1.32 H	IV	[<=0.89]	19-344-900041	10-Dec-19 09:57:00	10-Dec-19 09:57:00	10-Dec-19 09:59:15
HSV 1 Glycoprotein G Ab, IgG	3.56 H	IV	[<=0.89]	19-344-900041	10-Dec-19 09:57:00	10-Dec-19 09:57:00	10-Dec-19 10:00:10
HSV 2 Glycoprotein G Antibody, IgG	0.85	IV	[<=0.89]	19-344-900041	10-Dec-19 09:57:00	10-Dec-19 09:57:00	10-Dec-19 10:00:10

10-Dec-19 09:57:00 HSV Type 1/2 Combined Ab, IgG:

Specimen tested positive for Herpes Simplex Virus Type 1 and/or 2 Antibodies, IgG. ARUP test codes 0050292 and 0050294 will be added. Additional charges apply.

10-Dec-19 09:57:00 HSV Type 1/2 Combined Ab, IgG:  
 INTERPRETIVE INFORMATION: HSV 1/2 COMBINED Ab SCREEN, IgG  
 0.89 IV or less.....Not Detected  
 0.90-1.09 IV.....Indeterminate- Repeat testing  
   in 10-14 days may be helpful.  
 1.10 IV or greater.....Detected

The best evidence for current infection is a significant change on two appropriately timed specimens, where both tests are done in the same laboratory at the same time.

10-Dec-19 09:57:00 HSV 1 and/or 2 Abs, IgM by ELISA:  
 INTERPRETIVE INFORMATION: Herpes Simplex Virus Type 1 and/or 2 Antibodies, IgM by ELISA

0.89 IV or Less ..... Not Detected  
 0.90 - 1.09 IV ..... Indeterminate- Repeat testing in  
   10-14 days may be helpful.  
 1.10 IV or Greater ..... Detected-IgM antibody to HSV  
   detected, which may indicate a  
   current or recent infection.  
   However, low levels of IgM  
   antibodies may occasionally  
   persist for more than 12  
   months post-infection.

10-Dec-19 09:57:00 HSV 1 Glycoprotein G Ab, IgG:  
 REFERENCE INTERVAL: HSV 1 Glycoprotein G Ab, IgG

0.89 IV or less ..... Negative - No significant level of  
   detectable IgG antibody to HSV  
   type 1 glycoprotein G.  
 0.90 - 1.09 IV ..... Equivocal - Questionable presence  
   of IgG antibody to HSV type 1  
   glycoprotein G. Repeat testing in  
   10 - 14 days may be helpful.  
 1.10 IV or greater ... Positive - IgG antibody to HSV  
   type 1 glycoprotein G detected,  
   which may indicate a current or  
   past HSV infection.

\* Abnormal, # = Corrected, C = Critical, f = Footnote, H = High, L = Low, t = Interpretive Text, @ = Reference Lab

---

Individuals infected with HSV may not exhibit detectable IgG antibody to type-specific HSV antigens 1 and 2 in early stages of infection. Detection of antibody presence in these cases may only be possible using a non-type specific screening test.

10-Dec-19 09:57:00 HSV 2 Glycoprotein G Antibody, IgG:  
REFERENCE INTERVAL: HSV 2 Glycoprotein G Ab, IgG

- 0.89 IV or less ..... Negative - No significant level of detectable IgG antibody to HSV type 2 glycoprotein G.
- 0.90 - 1.09 IV ..... Equivocal - Questionable presence of IgG antibody to HSV type 2 glycoprotein G. Repeat testing in 10 - 14 days may be helpful.
- 1.10 IV or greater .... Positive - IgG antibody to HSV type 2 glycoprotein G detected, which may indicate a current or past HSV infection.

Individuals infected with HSV may not exhibit detectable IgG antibody to type-specific HSV antigens 1 and 2 in early stages of infection. Detection of antibody presence in these cases may only be possible using a non-type specific screening test.

\* Abnormal, # = Corrected, C = Critical, f = Footnote, H = High, L = Low, t = Interpretive Text, @ = Reference Lab