Patient Age/Gender: 49 years Male Printed: 09-Dec-19 09:55:53

Procedure	Result	Units	Ref Interval	Reported/ Accession Collected Received Verified
HSV 1 Glycoprotein G Ab, IgG	0.90 н	IV	[<=0.89]	19-340-900200 06-Dec-19 06-Dec-19 06-Dec-19 13:23:00 13:23:00 13:23:47
HSV 2 Glycoprotein G Antibody, IgG	0.90 н	IV	[<=0.89]	19-340-900200 06-Dec-19 06-Dec-19 06-Dec-19 13:23:00 13:23:00 13:23:47

06-Dec-19 13:23:00 HSV 1 Glycoprotein G Ab, IgG: REFERENCE INTERVAL: HSV 1 Glycoprotein G Ab, IgG

0.89 IV or less	Negative - No significant level of detectable IgG antibody to HSV
0.90 - 1.09 IV	type 1 glycoprotein G. Equivocal - Questionable presence
2107 21 11111	of IgG antibody to HSV type 1
	glycoprotein G. Repeat testing in 10 - 14 days may be helpful.
1.10 IV or greater	Positive - IgG antibody to HSV type 1 glycoprotein G detected, which may indicate a current or past HSV infection.

Individuals infected with HSV may not exhibit detectable IgG antibody to type-specific HSV antigens 1 and 2 in early stages of infection. Detection of antibody presence in these cases may only be possible using a non-type specific screening test.

06-Dec-19 13:23:00 HSV 2 Glycoprotein G Antibody, IgG: REFERENCE INTERVAL: HSV 2 Glycoprotein G Ab, IgG

0.89 IV or less	Negative - No significant level of detectable IgG antibody to HSV
0.90 - 1.09 IV	type 2 glycoprotein G. Equivocal - Questionable presence of IgG antibody to HSV type 2
	glycoprotein G. Repeat testing in 10 - 14 days may be helpful.
1.10 IV or greater	Positive - IgG antibody to HSV type 2 glycoprotein G detected, which may indicate a current or past HSV infection.

Individuals infected with HSV may not exhibit detectable IgG antibody to type-specific HSV antigens 1 and 2 in early stages of infection. Detection of antibody presence in these cases may only be possible using a non-type specific screening test.