



Procedure	Result	Units	Ref Interval	Accession	Collected	Received	Reported/Verified
HSV Type 1/2 Combined Ab, IgG	1.11 f	IV		19-343-900145	09-Dec-19 11:27:00	09-Dec-19 12:28:00	09-Dec-19 12:34:15
HSV 1 Glycoprotein G Ab, IgG	1.21 H	IV	[<=0.89]	19-343-900145	09-Dec-19 11:27:00	09-Dec-19 12:28:00	09-Dec-19 12:35:29
HSV 2 Glycoprotein G Antibody, IgG	0.91 H	IV	[<=0.89]	19-343-900145	09-Dec-19 11:27:00	09-Dec-19 12:28:00	09-Dec-19 12:35:29

09-Dec-19 11:27:00 HSV Type 1/2 Combined Ab, IgG:

Specimen tested positive for Herpes Simplex Virus Type 1 and/or 2 Antibodies, IgG. ARUP test codes 0050292 and 0050294 will be added. Additional charges apply.

09-Dec-19 11:27:00 HSV Type 1/2 Combined Ab, IgG:
 INTERPRETIVE INFORMATION: HSV 1/2 COMBINED Ab SCREEN, IgG
 0.89 IV or less.....Not Detected
 0.90-1.09 IV.....Indeterminate- Repeat testing
 in 10-14 days may be helpful.
 1.10 IV or greater.....Detected

The best evidence for current infection is a significant change on two appropriately timed specimens, where both tests are done in the same laboratory at the same time.

09-Dec-19 11:27:00 HSV 1 Glycoprotein G Ab, IgG:
 REFERENCE INTERVAL: HSV 1 Glycoprotein G Ab, IgG

0.89 IV or less Negative - No significant level of detectable IgG antibody to HSV type 1 glycoprotein G.
 0.90 - 1.09 IV Equivocal - Questionable presence of IgG antibody to HSV type 1 glycoprotein G. Repeat testing in 10 - 14 days may be helpful.
 1.10 IV or greater ... Positive - IgG antibody to HSV type 1 glycoprotein G detected, which may indicate a current or past HSV infection.

Individuals infected with HSV may not exhibit detectable IgG antibody to type-specific HSV antigens 1 and 2 in early stages of infection. Detection of antibody presence in these cases may only be possible using a non-type specific screening test.

09-Dec-19 11:27:00 HSV 2 Glycoprotein G Antibody, IgG:
 REFERENCE INTERVAL: HSV 2 Glycoprotein G Ab, IgG

0.89 IV or less Negative - No significant level of detectable IgG antibody to HSV type 2 glycoprotein G.
 0.90 - 1.09 IV Equivocal - Questionable presence of IgG antibody to HSV type 2 glycoprotein G. Repeat testing in 10 - 14 days may be helpful.
 1.10 IV or greater Positive - IgG antibody to HSV

* Abnormal, # = Corrected, C = Critical, f = Footnote, H = High, L = Low, t = Interpretive Text, @ = Reference Lab

type 2 glycoprotein G detected,
which may indicate a current or
past HSV infection.

Individuals infected with HSV may not exhibit detectable IgG antibody to type-specific HSV antigens 1 and 2 in early stages of infection. Detection of antibody presence in these cases may only be possible using a non-type specific screening test.