



Procedure	Result	Units	Ref Interval	Accession	Collected	Received	Reported/Verified
CASPR2 Ab IgG Screen by IFA, CSF	<b>Detected *</b>		[< 1:1]	19-260-900018	17-Sep-19 07:53:00	17-Sep-19 07:53:00	17-Sep-19 07:58:20
CASPR2 Ab IgG Titer by IFA, CSF	<b>1:40 *</b>		[< 1:1]	19-260-900018	17-Sep-19 07:53:00	17-Sep-19 07:53:00	17-Sep-19 07:58:25
LGI1 Ab IgG Screen by IFA, CSF	<b>Detected *</b>		[< 1:1]	19-260-900018	17-Sep-19 07:53:00	17-Sep-19 07:53:00	17-Sep-19 07:58:20
LGI1 Ab IgG Titer by IFA, CSF	<b>1:80 *</b>		[< 1:1]	19-260-900018	17-Sep-19 07:53:00	17-Sep-19 07:53:00	17-Sep-19 07:58:26
Voltage-Gated Potassium Channel Ab, CSF	<b>415.0 H</b>	pmol/L	[0.0-1.1]	19-260-900018	17-Sep-19 07:53:00	17-Sep-19 07:53:00	17-Sep-19 07:58:20

17-Sep-19 07:53:00 CASPR2 Ab IgG Screen by IFA, CSF  
 CASPR2 Antibody, IgG is detected. Titer results to follow.

17-Sep-19 07:53:00 LGI1 Ab IgG Screen by IFA, CSF  
 LGI1 Antibody, IgG is detected. Titer results to follow.

17-Sep-19 07:53:00 CASPR2 Ab IgG Screen by IFA, CSF:

INTERPRETIVE INFORMATION: CASPR2 Ab IgG w/Reflex  
 to Titer, CSF

Contactin-associated protein-2 (CASPR2) IgG antibody may occur as part of the voltage-gated potassium channel (VGKC) complex antibodies.

The presence of CASPR2 IgG antibody is associated with a wide spectrum of clinical manifestations, including acquired neuromyotonia, limbic encephalitis, painful neuropathy, and Morvan syndrome. Tumors such as thymoma, small cell lung cancer, and other rarer tumors may occur. The full-spectrum of clinical disorders and tumors associated with the CASPR2 IgG antibody continues to be defined. Results should be interpreted in correlation with the patient's clinical history and other laboratory findings.

This indirect fluorescent antibody assay utilizes contactin-associated protein-2 (CASPR2) transfected cell lines for the detection and semi-quantification of the CASPR2 IgG antibody.

Test developed and characteristics determined by ARUP Laboratories. See Compliance Statement D: aruplab.com/CS

17-Sep-19 07:53:00 CASPR2 Ab IgG Titer by IFA, CSF:

INTERPRETIVE INFORMATION: CASPR2 Ab Titer IgG by IFA, CSF

Test developed and characteristics determined by ARUP Laboratories. See Compliance Statement D: aruplab.com/CS

17-Sep-19 07:53:00 LGI1 Ab IgG Screen by IFA, CSF:

INTERPRETIVE INFORMATION: LGI1 Ab IgG w/Reflex to Titer, CSF

Leucine-rich, glioma-inactivated 1 protein (LGI1) IgG antibody may occur as part of the voltage-gated potassium channel (VGKC) complex antibodies.

The presence of LGI1 IgG antibody is mainly associated with limbic encephalitis, hyponatremia, and myoclonic movements. LGI1 IgG antibody is rarely associated with tumors

\* Abnormal, # = Corrected, C = Critical, f = Footnote, H = High, L = Low, t = Interpretive Text, @ = Reference Lab

---

but may occur infrequently in Morvan syndrome, neuromyotonia, and idiopathic epilepsy. The full-spectrum of clinical disorders associated with the LGI1 IgG antibody continues to be defined. Results should be interpreted in correlation with the patient's clinical history and other laboratory findings.

This indirect fluorescent antibody assay utilizes leucine-rich, glioma-inactivated 1 protein (LGI1) transfected cell lines for the detection and semi-quantification of the LGI1 IgG antibody.

Test developed and characteristics determined by ARUP Laboratories. See Compliance Statement D: aruplab.com/CS

17-Sep-19 07:53:00 LGI1 Ab IgG Titer by IFA, CSF:  
INTERPRETIVE INFORMATION: LGI1 Ab Titer IgG by IFA, CSF

Test developed and characteristics determined by ARUP Laboratories. See Compliance Statement D: aruplab.com/CS

17-Sep-19 07:53:00 Voltage-Gated Potassium Channel Ab, CSF:  
INTERPRETIVE INFORMATION: Voltage-Gated Potassium Channel  
(VGKC) Antibody, CSF

Voltage-Gated Potassium Channel (VGKC) antibodies are associated with neuromuscular weakness as found in neuromyotonia (also known as Issacs syndrome) and Morvan syndrome. VGKC antibodies are also associated with paraneoplastic neurological syndromes and limbic encephalitis; however, VGKC antibody-associated limbic encephalitis may be associated with antibodies to leucine-rich, glioma-inactivated 1 protein (LGI1) or contactin-associated protein-2 (CASPR2) instead of potassium channel antigens. A substantial number of VGKC-antibody positive cases are negative for LGI1 and CASPR2 IgG autoantibodies, not all VGKC complex antigens are known. The clinical significance of this test can only be determined in conjunction with the patient's clinical history and related laboratory testing.

Test developed and characteristics determined by ARUP Laboratories. See Compliance Statement D: aruplab.com/CS